

UMCES

UNIVERSITY OF MARYLAND CENTER for ENVIRONMENTAL SCIENCE
CHESAPEAKE BIOLOGICAL LABORATORY

MEASUREMENTS OF DENITRIFICATION IN AQUATIC ECOSYSTEMS:

LITERATURE REVIEW AND DATA REPORT

November 2005

Measurements of Denitrification in Aquatic Ecosystems: Literature Review and Data Report

Prepared by:

S. E. Greene¹, Graduate Research Assistant

**Prepared as a
companion volume for:**

Nitrogen in Estuaries
W.R. Boynton and W.M. Kemp
in:

Nitrogen in the Marine Environment
Edited by D.G. Capone, D.A. Bronk, M.R. Mulholland and E.J. Carpenter
(2006 in preparation)

November, 2005

University of Maryland System
Center for Environmental Science

¹Chesapeake Biological Laboratory
P.O. Box 38, Solomons, MD 20688-0038

TABLE OF CONTENTS

	page
Table of contents.....	ii
List of tables and figures.....	iii
Introduction.....	1
Microbial and ecological significance of denitrification.....	1
Measuring denitrification.....	1
Objectives.....	2
Methods.....	2
Summary of Preliminary Findings.....	6
Techniques used to measure denitrification.....	6
Denitrification in aquatic environments.....	7
Future Directions.....	9
References.....	10
Appendix A	
Measurements of denitrification in aquatic environments – data set.....	13
Appendix B	
References for data in Appendix A.....	15

LIST OF TABLES AND FIGURES

		page
Tables		
Table 1	Additional information reported with denitrification rates.	3
Table 2	Description of environmental groupings for denitrification measurement sites.	5
Table 3	Codes for techniques used to measure denitrification.	5
Table 4	Minimum, maximum, mean, median and range of denitrification rates measured in a variety of environments.	8
Figures		
Figure 1	Number of publications reporting denitrification rate measurements made with various techniques.	6
Figure 2	Frequency of denitrification rates measured in aquatic environments.	8
Figure 3	Frequency of denitrification rates measured in estuaries.	9

Introduction

With some exceptions, nitrogen (N) is the principal nutrient limiting primary production in coastal marine environments (Ryther and Dunstan 1971). Since the industrial revolution, fixation of biologically inaccessible N in the atmospheric pool to bioavailable forms in the terrestrial pool has doubled due to human activity (Vitousek et al. 1997). As a result, N inputs to the coastal ocean have increased. Elevated N concentrations in coastal waters support increased primary productivity, and the direct and indirect effects of enhanced production bring about readily observable, ecosystem-scale changes in coastal environments (e.g. Nixon 1995, Kemp et al. 2005). Recognition of the large impact of enhanced N inputs on coastal systems has fostered interest in removal mechanisms for N. Denitrification, the microbially-mediated conversion of dissolved nitrate (NO_3) to dinitrogen (N_2) gas, is an important pathway in the global N budget (Seitzinger 1988) as well as in the N budgets of many coastal ecosystems (Boynton et al. 1995). This process is of special interest in eutrophic coastal ecosystems because it results in permanent removal of anthropogenic N from aquatic systems.

Microbial and ecological significance of denitrification

Nitrate can be used by heterotrophic bacteria as an alternative terminal electron acceptor (TEA) in the absence of oxygen (O_2). Whether NO_3 or a different TEA is used depends on the specific redox regime of the environment, as well as the relative abundance of the various TEA's (Santschi et al. 1990). Bacterial use of NO_3 as a TEA involves a stepwise reduction to N_2 , with each step catalyzed by a different enzyme. Of particular interest, and perhaps of greatest ecological significance, is the step in which nitrite (NO_2) is reduced to nitric oxide (NO). At this point, N that is in a bioavailable, dissolved form becomes a biologically inert gas. Upon further reduction of NO to N_2 , the N reaches a more chemically stable state and diffuses to the atmosphere. This closes a major loop in the anthropogenic N cycle, as large amounts of N_2 are removed from the atmosphere by humans to make fertilizer. Nitrogen fixed from the atmospheric N pool for fertilizer is moved around the terrestrial landscape and applied to agricultural fields. Residual, biologically active forms of N may then be nitrified to NO_3 and move from the terrestrial N pool into inland and coastal waters (aquatic/oceanic N pool). Denitrifiers in aquatic and marine sediments return this N to the atmospheric pool as N_2 . In addition to returning N to the atmospheric pool, the ecological significance of denitrification is that it permanently removes from a system N that would otherwise be available for primary production. This sort of loss is of special interest in eutrophic, N-limited systems where substantial anthropogenic N inputs support high primary productivity, with a host of ecological and economic implications (Kemp et al. 2005). Denitrification is also of interest because it is the only mechanism (short of physical export) by which N is truly removed from the system; N taken up by biota or buried in sediments can be made available again via decomposition or erosion processes.

Measuring denitrification

A fundamental hurdle in measuring denitrification is that the increase of N_2 due to denitrification is small relative to atmospheric or dissolved aqueous background

concentrations of N_2 . Any direct measurement of denitrification must allow resolution of this small difference, and several techniques have been developed for this purpose. An earlier, indirect approach (the acetylene blockage technique) involved measurement of the less-abundant intermediate gas, N_2O , for which small changes in concentration are easier to resolve than for N_2 (e.g. Jorgensen and Soerensen 1985, Kemp et al. 1990). More recent approaches (e.g. isotopic techniques or use of $N_2:Ar$ ratios) involve direct measurement of N_2 (e.g. Nielsen and Glud 1996, Kana et al. 1998).

Objectives

This document is a review of denitrification measurements made during the past four-and-a-half decades in global aquatic environments. The primary objectives of this review are to:

1. Collect information regarding denitrification rates in different aquatic environments;
2. Characterize the distribution of denitrification rates that have been measured (i.e. maximum, minimum and median rates);
3. Document the use of different techniques for measuring denitrification, and the degree to which these have changed during the review period.

Methods

Literature reviewed in this document was located primarily within the Cambridge Scientific Abstracts (CSA) Aquatic Sciences and Fisheries Abstracts online database (www.csa.com). A search was conducted for all publications within the database containing the word “denitrification” in the abstract. Where journal-specific abstract or title databases were available, journals identified within the CSA database as containing at least one article with “denitrification” in the abstract were searched individually as well. Finally, literature cited in articles identified within the database was reviewed so as to detect publications not contained in the database. In a few cases, it was evident that measurements from a single study were reported in multiple publications, or previously-published data were included by another author in a table in a later publication. Care was taken to ensure that no measurement was reported more than once in this document. Several non-peer reviewed reports were also identified. Though it was not efficient to obtain copies of all of these works, it should be noted that denitrification measurements reported in gray literature are often published in peer reviewed venues as well. Due to the large number of measurements made with the acetylene blockage technique and the poor capacity of this technique to reflect natural conditions, some acetylene studies were judiciously rejected from inclusion in this review. It should also be noted that a substantial number of measurements of denitrification in terrestrial soils have been made (primarily using the acetylene blockage technique). An exhaustive review of that body of literature was not conducted, as the focus of this report is aquatic ecosystems.

In total, 164 studies were reviewed and 1929 measurements of denitrification recorded. The following environments were represented: inland (fresh) wetlands, freshwater creeks, lakes, coastal wetlands, mudflats, seagrass beds, lagoons, estuaries, reefs, continental shelves and human engineered systems (constructed wetlands and systems influenced directly by agriculture). The majority of reviewed measurements (60%) were made in estuarine systems. All measurements were converted to units of

$\mu\text{mol N m}^{-2} \text{ hr}^{-1}$ to facilitate comparisons. Where rates were originally reported on an areal basis, quantities of N were converted to μmoles , area converted to square meters and time to hours. Where rates were reported per volume of sediment or water, rates were multiplied by the sample depth or increment, based on information given in the publication.

Some rates were reported in units of N per gram of sediment weight. For these, information given in the text or assumptions regarding sediment bulk density were used for conversion to areal rates. Bulk densities of $0.2 - 0.9 \text{ g cm}^{-3}$ (derived after assumption of 60-80% water) were assumed for estuarine sediments, based on the equation:

$$\text{Bulk density (g cm}^{-3}\text{)} = \frac{(100 - \%H_2O)}{[(\%H_2O) + ((100 - \%H_2O)/2.5)]}$$

A bulk density of 1.4 was assumed for terrestrial soils.

In addition to denitrification rates, the following associated environmental parameters were recorded where data were available: geographic region, salinity class, temperature, month or season and year of collection, depth of measurement, station code or description, water column nitrate concentration, dissolved oxygen status of water adjacent to sediments, measurement technique, original units of measurement, and specific information regarding experimental manipulations (Table 1).

Table 1. Additional information reported with denitrification rates.

Column Title	Content Description
Location	Gives the name of the body of water or other environment in which denitrification measurements were made. In some cases, more information is given (e.g. specific region of a large body of water; country name; USA state name).
Environment Code	Classifies study areas by environment type (see Table 2).
Geographic Region	Gives the name of the general region in which studies were performed.
Station Code/Description	Gives the station name or code given by authors for specific sites in which measurements were made; or gives important information about site characteristics (e.g. high marsh, low marsh, algal mat, sediment characteristics).
Salinity Class	Gives the salinity class where measurements were made (Hypersaline, >40; Euhaline, 30-40; Polyhaline, 18-30; Mesohaline, 5-18; Oligohaline, 0.5-5; Tidal Fresh, <0.5 with tidal influence; Nontidal Fresh, <0.5 without tidal influence).
Station or Sample Depth	Gives the depth of stations where sediment denitrification measurements were made, or the depth at which water column denitrification measurements were made (numbers indicating depths of water column samples are followed by an asterisk "**").

Table 1 cntnd.

Column Title	Content Description
Year	Gives the year in which measurements were made.
Month/Season	Gives the month or season in which measurements were made.
Temperature	Gives the water temperature at which measurements were made.
Nitrate	Gives the nitrate (NO ₃) concentration from the water column. In some cases, ambient NO ₃ concentration was not specified, but the amount added for experimental purposes was (identified by an asterisk "*").
Oxygen	Gives the oxygen (O ₂) concentration from the water column. In several publications, only general information describing aerobic, anaerobic, hypoxic or anoxic conditions was given. Such conditions were included in this report in lieu of actual concentrations.
Denitrification	Gives the denitrification rate measured, with all rates converted to $\mu\text{mol N m}^{-2} \text{hr}^{-1}$. When direct and coupled denitrification were measured separately, both rates are given. When a range of rates was measured, gives the full range.
Simplified Denitrification	Gives the total denitrification rate measured, in $\mu\text{mol N m}^{-2} \text{hr}^{-1}$, with direct and coupled denitrification rates summed and expressed as a single rate and ranges of rates expressed as a mean rate.
Method Description	Describes (briefly) the technique used to measure denitrification.
Technique code	Classifies techniques into general categories. (See Table 3)
Reference	Citations for publications in which measurements were reported.
Unit Conversions	Gives information as to whether or not a unit conversion was necessary to express published denitrification rates as $\mu\text{mol N m}^{-2} \text{hr}^{-1}$, as well as the original, published units. In several cases, it was unclear whether published rates were in units of "N", "N ₂ ," or "N ₂ O". If the authors could not be contacted for clarification, assumptions are stated.
Comments	Gives additional information on specific experimental conditions/manipulations, whether or not rates had to be estimated from a graph, and in some cases important conclusions made by the authors.
Original Values	When a unit conversion was made for which the conversion process was not immediately obvious, original published rates and any assumptions necessary to make the conversion are listed.
Assumptions	See "Original Values" above.

Table 2. Description of environmental groupings for denitrification measurement sites.

Environment	Code	Description
Human engineered	H	Systems constructed or heavily impacted by humans (e.g. aquaculture, constructed wetlands, immediate agricultural runoff)
Wetland (fresh)	W	Landlocked wetland receiving only freshwater inputs
Creek (fresh)	Ck	Natural stream of water smaller than a river; labeled as such by the author(s)
Lake	Lk	Landlocked bodies of standing, fresh water
Coastal wetland	CW	Wetlands receiving tidal inputs (i.e. tidal marshes and mangroves)
Mudflat	MF	Unvegetated bar of sediment submerged regularly by tides
Seagrass Bed	SB	Area densely populated by submerged aquatic vegetation; labeled as such by the authors
Lagoon	Lg	Shallow body of water, cut off from open coastal waters by sand bars; labeled as such by the authors
Estuary	E	Semi-enclosed bodies of water receiving inputs from both terrestrial (freshwater) sources and the ocean
Reef	R	Sites in close proximity to coral reefs
Continental Shelf	CS	Open coastal waters near a major landmass

Table 3. Codes for various techniques used to measure denitrification.

Code	Technique
A	Acetylene
ISA	<i>In situ</i> acetylene
DN2	Direct measurement of N ₂ flux
ISDN2	<i>In situ</i> direct measurement of N ₂ flux
GP	Gas Partitioning
HA	"Heavy" ammonium (¹⁵ NH ₄) addition
HN	"Heavy" nitrate (¹⁵ NO ₃) addition
HNA	"Heavy" nitrate & ammonium addition
IP	Isotope Pairing
M	MIMS
MB	Mass balance
MO	Miscellaneous other

Summary of Preliminary Findings

Techniques used to measure denitrification

Scientists have been aware of denitrification for over a century (Zumft 1997). Though investigations of denitrification were published in the 1950's (e.g. Wijler and Delwiche 1954, Hauck et al. 1958), attempts to actually *measure* the process were not made until the 1960's (Goering and Dugdale 1966). The earliest estimates of denitrification were made via N mass balances and isotopic studies in which $^{15}\text{N-NO}_3$ was added to an environment and dissolved N_2 was analyzed for ^{15}N after an incubation period (Goering and Dugdale 1966). Important new measurement techniques have appeared since the 1960's, most notably the acetylene blockage technique in the 1970's (e.g. Balderston et al. 1976), direct measurement of N_2 production from sediment cores in the 1980's (e.g. Seitzinger et al. 1980, Seitzinger 1988), and the MIMS (membrane inlet mass spectrometry, Kana et al. 1994) and isotope pairing (Nielsen 1992) techniques in the 1990's.

More denitrification measurements appear to have been made with the acetylene blockage technique than with any other technique (Fig. 1). This technique owes its popularity to the fact that it circumvents the central quandary in measuring denitrification - the need to distinguish relatively small changes in N_2 against a large atmospheric (and by equilibration, dissolved aqueous) background. By blocking the final step in denitrification (reduction of N_2O to N_2), acetylene permits accumulation and detection of the more easily measurable gas, N_2O , as a proxy for N_2 (Balderston et al. 1976). However, acetylene also inhibits nitrification (Hynes and Knowles 1978), which can be an important source of NO_3 for denitrification, so this technique can lead to underestimates of *in situ* rates (Kemp et al. 1990). Acetylene studies do not appear as popular in the current decade as in previous decades, likely due to the development of other direct techniques that do not involve inhibitors and associated problems (Fig. 1).




Figure 1. Number of publications reporting denitrification rate measurements made with various techniques.

Seitzinger made direct measurements of N_2 fluxes by lowering the background N_2 concentration via sparging core headspaces with an atmosphere-like mixture of gases in which N_2 is replaced by helium (Seitzinger et al. 1980). This technique has the disadvantage of requiring long pre-incubation periods, and despite attempts at correction, there is also the issue of distinguishing the N_2 flux due to denitrification from the flux due to chemical gradients (Seitzinger 1988, Cornwell et al. 1999). Nowicki (1994) has used anoxic control cores to overcome the problem of sediment-porewater degassing, such that the anoxic core flux of N_2 can be assumed to be due solely to chemical gradients. The reducing conditions brought on by anoxia cause quick depletion of NO_3 stocks and also inhibit nitrification, both of which prevent denitrification from occurring in anoxic control cores.

A large number of denitrification measurements have been made with the isotope pairing technique since the 1990's (e.g. Risgaard-Petersen et al. 1994, Rysgaard et al. 1995, Trimmer et al. 1999, Dong et al. 2000). In this method, $^{15}N-NO_3$ is added to overlying water and N_2 evolved during incubation is analyzed for partitioning into $^{15}N^{15}N$, $^{15}N^{14}N$ and $^{14}N^{14}N$ (Nielsen 1992). The advantage of this technique is that the different pathways for denitrification can be distinguished (direct denitrification of water column NO_3 versus coupled nitrification-denitrification). By 2004 more denitrification measurements had been made with isotope pairing than with any other technique in the current decade (Fig. 1). However, several (perhaps questionable) underlying assumptions are required, namely that added $^{15}NO_3$ mixes homogeneously with the $^{14}NO_3$ pool, that added NO_3 does not change the rate of coupled nitrification-denitrification, that isotope fractionation can be neglected, and that diffusion of heavy and light NO_3 is similar (Middleburg et al. 1996).

In the mid-1990's, the MIMS technique was first applied to the study of denitrification in estuarine sediments (Kana et al. 1994, Kana et al. 1998). A membrane inlet mass spectrometer optimized for high-precision measurement of dissolved gasses is employed in this technique. A measurement coefficient of variation of <0.05% is achieved by measuring N_2 to argon (Ar) ratios in water samples, compared to precision levels of 0.5% when measuring N_2 alone (Kana et al. 1994). This technique has gained popularity since its inception, though it has incited criticism from proponents of the isotope pairing technique (Eyre et al. 2002, Eyre et al. 2004). Even so, MIMS remains one of the preferred techniques for measuring denitrification in aquatic ecosystems. In general, the study of denitrification has gained momentum, and nearly as many studies using isotope pairing and MIMS have already been published in the first few years of the present decade as were published in the entire previous decade (Fig. 1).

Denitrification in aquatic environments

Since the 1960's, denitrification has been measured in a large variety of environments, both natural and engineered. Denitrification rates between 10 and 100 $\mu\text{moles N m}^{-2} \text{h}^{-1}$ are most frequently reported, though rates well over 1000 $\mu\text{moles N m}^{-2} \text{h}^{-1}$ have been observed in certain systems (Fig. 2; Table 4). Rates near 20,000 $\mu\text{moles N m}^{-2} \text{h}^{-1}$ have been measured in human engineered systems and estuaries, though median rates for these environments are not so high as to be outside the range of commonly measured rates (Table 4; Fig. 2). Median rates appear highest in lakes (37 $\mu\text{mol N m}^{-2} \text{h}^{-1}$), estuaries (40 $\mu\text{mol N m}^{-2} \text{h}^{-1}$) and coastal wetlands (54 $\mu\text{mol N m}^{-2} \text{h}^{-1}$; Table 4). In




Figure 2. Frequency distribution of denitrification rates measured in aquatic environments.

estuaries, where more measurements of denitrification appear to have been made than in any other environment (Table 4), the most commonly reported rates are between 1 and 100 $\mu\text{mol N m}^{-2} \text{h}^{-1}$ (Fig. 3), and more specifically between 1 and 50 $\mu\text{mol N m}^{-2} \text{h}^{-1}$. The range of rates measured in estuaries is higher than that of any other natural system by an order of magnitude (Table 4).

Table 4. Minimum, maximum, mean, median and range of denitrification rates measured in a variety of aquatic environments. “n” = number of measurements. Rates are given in $\mu\text{mol N m}^{-2} \text{h}^{-1}$.

	Min	Max	Mean	Median	Range	n
Human engineered	0	24143	695	1	24143	68
Wetlands (fresh)	0	330	39	4	330	52
Creeks (fresh)	0	1200	195	32	1200	20
Lake	0	490	89	37	490	90
Coastal wetland	-200	1865	94	54	2065	167
Mudflat	2	213	71	31	211	61
Seagrass Bed	2	167	29	8	165	13
Lagoon	-20	290	21	9	310	116
Estuary	-93	19616	197	40	19709	1188
Reef	0	533	58	4	533	40
Continental Shelf	0	1658	104	15	1658	113




Figure 3. Frequency distribution of denitrification rates measured in estuaries.

Future Directions

Synthetic analyses of existing data promote scientific efficiency by highlighting needs for additional measurements and, perhaps more importantly, revealing areas for which additional measurement efforts would be redundant. To that end, data from this review should facilitate analyses of relationships between denitrification rate and environmental parameters such as salinity, temperature, oxygen concentration and depth. Such analyses will further a mechanistic understanding of denitrification and may aid in modeling efforts. These data may also be used to examine distributions of denitrification rates within specific environments (e.g. Fig. 3). Due to the large amount of information, all data are located in an electronic appendix (Microsoft Excel file “Appendix A.xls”) on the accompanying CD.

References

- Balderston, W. L., B. Sherr and W.J. Payne. 1976. Blockage by acetylene of nitrous oxide reduction in *Pseudomonas perfectomarinus*. Applied and Environmental Microbiology **31**: 504-508.
- Boynton, W. R., J.H. Garber, R. Summers and W.M Kemp. 1995. Inputs, transformations and transport of nitrogen and phosphorus in Chesapeake Bay and selected tributaries. Estuaries **18**: 285-314.
- Cornwell, J. C., W.M. Kemp and T.M. Kana. 1999. Denitrification in coastal ecosystems: methods, environmental controls, and ecosystem level controls, a review. Aquatic Ecology **33**: 41-54.
- Dong, L. F., D.C. Thornton, D.B. Nedwell and G.J. Underwood. 2000. Denitrification in sediments of the River Colne estuary, England. Marine Ecology Progress Series **203**: 109-122.
- Eyre, B. D., S. Rysgaard, T. Dalsgaard and P.B. Christensen. 2002. Comparison of isotope pairing and N₂:Ar methods for measuring sediment denitrification - assumptions, modifications, and implications. Estuaries **25**: 1077-1087.
- . 2004. Reply to comment on our paper "Comparison of isotope pairing and N₂:Ar methods for measuring sediment denitrification". Estuaries **27**: 177-178.
- Goering, J. J. and V. A. Dugdale. 1966. Estimates of the rates of denitrification in a subarctic lake. Limnology & Oceanography **11**: 113-117.
- Hauck, R. D., S.W. Melsted and P.E. Yankwich. 1958. Use of N-Isotope distribution in nitrogen gas in the study of denitrification. Soil Science **86**: 287-291.
- Hynes, R. K. and R. Knowles. 1978. Inhibition by acetylene of ammonia oxidation in *Nitrosomonas europaea*. FEMS Microbial Letters **4**: 319-321.
- Jorgensen, B. B. and J. Sorensen. 1985. Seasonal cycles of O₂, NO₃ and SO₄ reduction in estuarine sediments: the significance of a NO₃ reduction maximum in spring. Marine Ecology Progress Series **24**: 65-74.
- Kana, T. M., C. Darkangelo, M.D. Hunt, J.B. Oldham, G.E. Bennett and J.C. Cornwell. 1994. Membrane Inlet Mass Spectrometer for rapid high-precision determination of N₂, O₂ and Ar in environmental water samples. Analytical Chemistry **66**: 4166-4170.
- Kana, S., J. Cornwell and K. Groszkowski. 1998. Denitrification in estuarine sediments determined by membrane inlet mass spectrometry. Limnology & Oceanography **43**: 334-339.

- Kemp, W. M., P.Sampou, J.Caffrey, M.Mayer, K. Henriksen and W.R. Boynton. 1990. Ammonium recycling versus denitrification in Chesapeake Bay sediments. *Limnology & Oceanography* **35**: 1545-1563.
- Kemp, W. M., W. Boynton, J. Adolf, D. Boesch, W. Boicourt, G. Brush, J. Cornwell, T. Fisher, P. Glibert, J. Hagy, L. Harding, E. Houde, D. Kimmel, W. Miller, R. Newell, M. Roman, E. Smith and J.C. Stevenson. 2005. Eutrophication of Chesapeake Bay: historical trends and ecological interactions. *Marine Ecology Progress Series* *in press*.
- Middleburg, J. J., K.Soetaert and P.M. Herman. 1996. Evaluation of the nitrogen isotope-pairing method for measuring benthic denitrification: A simulation analysis. *Limnology & Oceanography* **41**: 1839-1844.
- Nielsen, L. P. 1992. Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbial Ecology* **86**: 357-362.
- Nielsen, L. P. and R. Glud. 1996. Denitrification in a coastal sediment measured in situ by the nitrogen isotope pairing technique applied to a benthic flux chamber. *Marine Ecology Progress Series* **137**: 181-186.
- Nixon, S. W. 1995. Coastal marine eutrophication: A definition, social causes, and future concerns. *Ophelia* **41**: 199-219.
- Nowicki, B. L. 1994. The effect of temperature, oxygen, salinity, and nutrient enrichment on estuarine denitrification rates measured with a modified nitrogen gas flux technique. *Estuarine, Coastal and Shelf Science* **38**: 137-156.
- Risgaard-Petersen, N., S. Rysgaard, L.P. Nielsen and N.P. Revsbech. 1994. Diurnal variation of denitrification and nitrification in sediments colonized by benthic microphytes. *Limnology & Oceanography* **39**: 573-579.
- Rysgaard, S., P.B. Christensen and L.P. Nielsen. 1995. Seasonal variation in nitrification and denitrification in estuarine sediment colonized by benthic microalgae and bioturbating infauna. *Marine Ecology Progress Series* **126**: 111-121.
- Ryther, J. H. and W. M. Dunstan. 1971. Nitrogen, phosphorus, and eutrophication in the coastal marine environment. *Science* **171**: 1008-1013.
- Santschi, P., P. Hoehener, G. Benoit and M. Buchholtz-Ten Brink. 1990. Chemical processes at the sediment-water interface. *Marine Chemistry* **30**: 269-315.
- Seitzinger, S. P., S.W. Nixon, M.E. Pilson and S. Burke. 1980. Denitrification and N₂O production in nearshore marine sediments. *Geochim. Cosmochim. Acta* **44**: 1853-1860.

- Seitzinger, S. P. 1988. Denitrification in freshwater and coastal marine ecosystems: Ecological and geochemical significance. *Limnology & Oceanography* **33**: 702-724.
- Trimmer, M., R.J. Gowen, B.M. Stewart and D.B. Nedwell. 1999. The spring bloom and its impact on benthic mineralisation rates in western Irish Sea sediments. *Marine Ecology Progress Series* **185**: 37-46.
- Vitousek, P. M., J.D. Aber, R.W. Howarth, G.E. Likens, P.A. Matson, D.W. Schindler, W.H. Schlesinger and D.G. Tilman. 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecological Applications* **7**: 737-750.
- Wijler, J. and C.C. Delwiche. 1954. Investigations on the denitrifying process in soil. *Plant Soil* **5**: 155-169.
- Zumft, W. G. 1997. Cell biology and molecular basis of denitrification. *Microbiology and Molecular Biology Reviews* **61**: 533-616.

Appendix A

*Measurements of Denitrification in
Aquatic Environments – Data Set*

Appendix B

*References for Data in
Appendix A*

References

- Abd Aziz, S. A. and D. B. Nedwell. 1986. The nitrogen cycle of an east coast, U.K. salt marsh. 2. Nitrogen fixation, nitrification, denitrification, tidal exchange. *Estuarine, Coastal and Shelf Science* **22**: 689-704.
- Abril, G., S. Riou, H. Etcheber, M. Frankignoulle, R. De Wit and J. Middelburg. 2000. Transient, tidal time-scale, nitrogen transformations in an estuarine turbidity maximum-fluid mud system (The Gironde, South-west France). *Estuarine, Coastal and Shelf Science* **50**: 703-715.
- Ahlgren, I. 1967. Limnological studies of Lake Norrviken, a eutrophicated Swedish lake. 1. Water chemistry and nutrient budget. *Hydrologie* **29**: 53-90.
- Aller, R. C., J.E. Mackin, W.J. Ullman, W. Chen-Hou, T. Shing-Min, J. Jian-Cai, S. Yong-Nian and H. Jia-Zhen. 1985. Early chemical diagenesis, sediment-water solute exchange, and storage of reactive organic matter near the mouth of the Changjiang, East China Sea. *Continental Shelf Research* **4**: 227-251.
- Alongi, D., F. Tirendi, L. Trott and T. Xuan. 2000. Benthic decomposition rates and pathways in plantations of the mangrove *Rhizophora apiculata* in the Mekong delta, Vietnam. *Marine Ecology Progress Series* **194**: 87-101.
- An, S. and W. Gardner. 2002. Dissimilatory nitrate reduction to ammonium (DNRA) as a nitrogen link, versus denitrification as a sink in a shallow estuary (Laguna Madre /Baffin Bay, Texas). *Marine Ecology Progress Series* **237**: 41-50.
- Andersen, J. M. 1971. Nitrogen and phosphorus budgets and the role of sediments in six shallow Danish lakes. *Arch. Hydrobiol.* **74**: 528-550.
- . 1977. Rates of denitrification of undisturbed sediment from six lakes as a function of nitrate concentration, oxygen and temperature. *Arch. Hydrobiol.* **80**: 147-159.
- Anderson, I., C. Tobias, B. Neikirk and R. Wetzel. 1997. Development of a process-based nitrogen mass balance model for a Virginia (USA) *Spartina alterniflora* salt marsh: Implications for net DIN flux. *Marine Ecology Progress Series* **159**: 13-27.
- Bianchi, M., P. Bonin and Feliatra. 1994. Bacterial nitrification and denitrification rates in the Rhone River plume (northwestern Mediterranean Sea). *Marine Ecology Progress Series* **103**: 197-202.
- Billen, G. 1978. A budget of nitrogen recycling in North Sea sediments off the Belgian coast. *Estuarine Coastal Marine Science* **7**: 127-146.

- Blackburn, T. H., B.A. Lund and M.D. Krom. 1988. C- and N-mineralization in the sediments of earthen marine fishponds. *Marine Ecology Progress Series* **44**: 221-227.
- Blackburn, T. H., D.B. Nedwell and W.J. Wiebe. 1994. Active mineral cycling in a Jamaican seagrass sediment. *Marine Ecology Progress Series* **110**: 233-239.
- Blackburn, T. H., P. Hall, S. Hulth and A. Landen. 1996. Organic-N loss by efflux and burial associated with a low efflux of inorganic N and with nitrate assimilation in Arctic sediments (Svalbard, Norway). *Marine Ecology Progress Series* **141**: 283-293.
- Boynton, W. R., J.H. Garber, R. Summers, and W.M Kemp. 1995. Inputs, transformations and transport of nitrogen and phosphorus in Chesapeake Bay and selected tributaries. *Estuaries* **18**: 285-314.
- Brettar, I. and G. Rheinheimer. 1992. Influence of carbon availability on denitrification in the central Baltic Sea. *Limnology & Oceanography* **37**: 1146-1163.
- Burford, M. A. and A. R. Longmore. 2001. High ammonium production from sediments in hypereutrophic shrimp ponds. *Marine Ecology* **224**: 187-195.
- Cabrita, M. T. and V. Brotas. 2000. Seasonal variation in denitrification and dissolved nitrogen fluxes in intertidal sediments of the Tagus Estuary, Portugal. *Marine Ecology Progress Series* **202**: 51-65.
- Caffrey, J. M. and W. M. Kemp. 1990. Nitrogen cycling in sediments with estuarine populations of *Potamogeton perfoliatus* and *Zostera marina*. *Marine Ecology Progress Series* **66**: 147-160.
- . 1992. Influence of the submersed plant, *Potamogeton perfoliatus* , on nitrogen cycling in estuarine sediments. *Limnology & Oceanography* **37**: 1483-1495.
- Capone, D., S.E. Dunham, S.G. Horrigan and L.E. Duguay. 1992. Microbial nitrogen transformations in unconsolidated coral reef sediments. *Marine Ecology Progress Series* **80**: 75-88.
- Chan, Y. K. and N. R. Campbell. 1980. Denitrification in Lake 227 during summer stratification. *Can. J. Fish Aquatic. Sci.* **37**: 506-512.
- Childs, C. R., N.N. Rabalais, R.E. Turner and L.M. Proctor. 2002. Sediment denitrification in the Gulf of Mexico zone of hypoxia. *Marine Ecology Progress Series* **240**: 285-290.

- Christensen, J. P., W.H. Smethie and A.H. Devol. 1987a. Benthic nutrient regeneration and denitrification on the Washington continental shelf. *Deep-Sea Research* **34**: 1027–1049.
- Christensen, J. P., J.W. Murray, A.H. Devol and L.A. Codispoti. 1987b. Denitrification in continental shelf sediments has major impact on the oceanic nitrogen budget. *Global Biogeochemical Cycles* **1**: 97-116.
- Christensen, P. B., L.P. Nielsen, J. Sorensen, N.P. Revsbech. 1990. Denitrification in nitrate-rich streams: diurnal and seasonal variation related to benthic oxygen metabolism. *Limnology & Oceanography* **35**: 640-651.
- Christensen, P. B. and J. Sorensen. 1986. Temporal variation of denitrification activity in plant-covered littoral sediment from Lake Hampen, Denmark. *Applied and Environmental Microbiology* **51**: 1174-1179.
- Cooper, A. B. 1990. Nitrate depletion in the riparian zone and stream channel of a small headwater catchment. *Hydrobiologia* **202**: 13-26.
- Currin, C. A., S.B. Joye, and H.W. Paerl. 1996. Diel rates of N sub(2)-fixation and denitrification in a transplanted *Spartina alterniflora* marsh: Implications for N-flux dynamics. *Estuarine, Coastal and Shelf Science* **42**: 597-616.
- Debusk, W. F. and K. R. Reddy. 1987. Removal of floodwater nitrogen in a cypress swamp receiving primary wastewater effluent. *Hydrobiologia* **153**: 79-86.
- Devol, A. H. 1991. Direct measurement of nitrogen gas fluxes from continental shelf sediments. *Nature* **349**: 319-321.
- Devol, A. H. and J. P. Christensen. 1993. Benthic fluxes and nitrogen cycling in sediments of the continental margin of the eastern North Pacific. *Journal of Marine Research* **51**: 345-372.
- Dong, L. F., D.C. Thornton, D.B. Nedwell, and G.J. Underwood. 2000. Denitrification in sediments of the River Colne estuary, England. *Marine Ecology Progress Series* **203**: 109-122.
- Duff, J. H., F.J. Triska and R.S. Oremland. 1984. Denitrification associated with stream periphyton: chamber estimates from undisrupted communities. *Journal of Environmental Quality* **13**: 514-518.
- Eriksson, P. G. and S. B. Weisner. 1997. Nitrogen removal in a wastewater reservoir: The importance of denitrification by epiphytic biofilms on submersed vegetation. *Journal of Environmental Quality* **26**: 905-910.

- . 1999. An experimental study on effects of submersed macrophytes on nitrification and denitrification in ammonium-rich aquatic systems. *Limnology & Oceanography* **44**: 1993-1999.
- Eyre, B. D., S. Rysgaard, T. Dalsgaard and P.B. Christensen. 2002. Comparison of isotope pairing and N₂:Ar methods for measuring sediment denitrification - assumptions, modifications, and implications. *Estuaries* **25**: 1077-1087.
- Eyre, B. D. and A. J. Ferguson. 2002. Comparison of carbon production and decomposition, benthic nutrient fluxes and denitrification in seagrass, phytoplankton, benthic microalgae and macroalgae-dominated warm-temperate Australian lagoons. *Marine Ecology Progress Series* **229**: 43-59.
- Garcia-Ruiz, R., S. Pattinson and B. Whitton. 1998. Denitrification in river sediments: relationship between process rate and properties of water and sediment. *Freshwater Biology* **39**: 467-476.
- Gardner, W. S., T.F. Nalepa and J.M. Malczyk. 1987. Nitrogen mineralization and denitrification in Lake Michigan sediments. *Limnology & Oceanography* **32**: 1226-1238.
- Giblin, A. E., C. Hopkinson and J. Tucker. 1993. Metabolism, nutrient cycling and denitrification in Boston Harbor and Massachusetts Bay sediments. Massachusetts Water Resource Authority. Environmental Quality Department Technical Report 93-2. 46 p.
- Gilbert, F., P. Souchu, M. Bianchi and P. Bonin. 1997. Influence of shellfish farming activities on nitrification, nitrate reduction to ammonium and denitrification at the water-sediment interface of the Thau Lagoon, France. *Marine Ecology Progress Series* **151**: 143-153.
- Gilbert, F., G. Stora, P. Bonin and K. Aarnio. 1998. Influence of bioturbation on denitrification activity in Mediterranean coastal sediments: an in situ experimental approach. *Marine Ecology Progress Series* **163**: 99-107.
- Glud, R. N., O. Holby, F. Hoffmann and D.E. Canfield. 1998. Benthic mineralization and exchange in Arctic sediments (Svalbard, Norway). *Marine Ecology Progress Series* **173**: 237-251.
- Glud, R. N., N. Risgaard-Petersen, B. Thamdrup, H. Fossing and S. Rysgaard. 2000. Benthic carbon mineralization in a high-Arctic sound (Young Sound, NE Greenland). *Marine Ecology Progress Series* **206**: 59-71.
- Goering, J. J. and V. A. Dugdale. 1966. Estimates of the rates of denitrification in a subarctic lake. *Limnology & Oceanography* **11**: 113-117.

- Griffiths, R. P., B.A. Caldwell, W.A. Broich and R.Y. 1982. The long-term effects of crude oil on microbial processes in subarctic marine sediments. *Estuarine, Coastal and Shelf Science* **15**: 183-198.
- Groffman, P. M., E.A. Axelrod, J.L. Lemunyon and W.M. Sullivan. 1991. Denitrification in grass and forest vegetated filter strips. *Journal of Environmental Quality* **20**: 671-674.
- Haines, E. B., A. Chalmers, R. Hanson and B. Sherr. 1977. Nitrogen pools and fluxes on a Georgia salt marsh. *In* M. Wiley [ed.], *Estuarine Processes*. Academic Press.
- Haines, J. R., R.M. Atlas, R.P. Griffiths and R.Y. Morita. 1981. Denitrification and nitrogen fixation in Alaskan continental shelf sediments. *Applied and Environmental Microbiology* **41**: 412-421.
- Hall, P. J., O. Holby, S. Kollberg and M-O. Samuelsson. 1992. Chemical fluxes and mass balances in a marine fish cage farm. 4. Nitrogen. *Marine Ecology Progress Series* **89**: 81-91.
- Hemminga, M., P. Harrison and F. Van Lent. 1991. The balance of nutrient losses and gains in seagrass meadows. *Marine Ecology Progress Series* **71**: 85-96.
- Hemond, H. F. 1983. The nitrogen budget of Thoreau's bog. *Ecology* **64**: 99-109.
- Henrikson, K. and W. M. Kemp. 1988. Nitrification in estuarine and coastal marine sediments, p. 207-249. *In* T. H. a. J. S. Blackburn [ed.], *Nitrogen cycling in coastal marine environments*. Proc. SCOPE Symp. 33. Wiley.
- Hietanen, S., P.H. Moisander, J. Kuparinen and L. Tuominen. 2002. No sign of denitrification in a Baltic Sea cyanobacterial bloom. *Marine Ecology Progress Series* **242**: 73-82.
- Hines, M. E., D.A. Bazylnski, J.B. Tugel and W.B. Lyons. 1991. Anaerobic microbial biogeochemistry in sediments from two basins in the Gulf of Maine: Evidence for iron and manganese reduction. *Estuarine, Coastal and Shelf Science* **32**: 313-324.
- Holmes, R. M., J.B. Jr. Jones, S.G. Fisher and N.B. Grimm. 1996. Denitrification in a nitrogen-limited stream ecosystem. *Biogeochemistry* **33**: 125-146.
- Iizumi, H., A. Hattori and C.P. Mcroy. 1980. Nitrate and nitrite in in interstitial waters of eelgrass beds in relation to the rhizosphere. *J. Exp. Mar. Biol. Ecol.* **47**: 191-201.
- Jenkins, M. C. and W. M. Kemp. 1984. The coupling of nitrification and denitrification in two estuarine sediments. *Limnology & Oceanography* **29**: 609-619.

- Jensen, K. M., M.H. Jensen and E. Kristensen. 1996. Nitrification and denitrification in Wadden Sea sediments (Königshafen, Island of Sylt, Germany) as measured by nitrogen isotope pairing and isotope dilution. *Aquatic Microbial Ecology* **11**: 181–191.
- Jensen, M. H., T.K. Andersen and J. Soerensen. 1988. Denitrification in coastal bay sediment: Regional and seasonal variation in Aarhus Bight, Denmark. *Marine Ecology Progress Series* **48**: 155-162.
- Johnson, R., M. Lamontagne and I. Valiela. 1994. Rate of denitrification in submerged slat marsh sediments. *Biological Bulletin* **187**: 289-290.
- Jorgensen, B. B. and J. Sorensen. 1985. Seasonal cycles of O₂, NO₃⁻ and SO₄²⁻ reduction in estuarine sediments: the significance of a NO₃⁻ reduction maximum in spring. *Marine Ecology Progress Series* **24**: 65-74.
- . 1988. Two annual maxima of nitrate reduction and denitrification in estuarine sediment (Norsminde Fjord, Denmark). *Marine Ecology Progress Series* **48**: 147-154.
- Joye, S. B., S.V. Smith, J.T. Hollibaugh and H.W. Paerl. 1996. Estimating denitrification rates in estuarine sediments: A comparison of stoichiometric and acetylene based methods. *Biogeochemistry* **33**: 197-215.
- Joye, S. B. and H. W. Paerl. 1993. Contemporaneous nitrogen fixation and denitrification in intertidal microbial mats: Rapid response to runoff events. *Marine Ecology Progress Series* **94**: 267-274.
- . 1994. Nitrogen cycling in microbial mats: Rates and patterns of denitrification and nitrogen fixation. *Marine Biology* **119**: 285-295.
- Kana, T., Sullivan, J.C. Cornwell, and K. Groszkowski. 1998. Denitrification in estuarine sediments determined by membrane inlet mass spectrometry. *Limnology & Oceanography* **43**: 334-339.
- Kaplan, W., I. Valiela and J.M. Teal. 1979. Denitrification in a salt marsh ecosystem. *Limnology & Oceanography* **24**: 726-734.
- Kaspar, H. F. 1983. Denitrification, nitrate reduction to ammonium and inorganic nitrogen pools in intertidal sediments. *Marine Biology* **74**: 133-139.
- Kaspar, H. F., R.A. Asher and I.C. Boyer. 1985a. Microbial nitrogen transformations in sediments and inorganic nitrogen fluxes across the sediment/water interface on the South Island west coast, New Zealand. *Estuarine, Coastal and Shelf Science* **21**: 245-255.

- Kaspar, H. F., P.A. Gillespie, I.C. Boyer and A.L. Mackenzie. 1985b. Effects of mussel aquaculture on the nitrogen cycle and benthic communities in Kenepuru Sound, Marlborough Sounds, New Zealand. *Marine Biology* **85**: 127-136.
- Keeney, D. R., R.L. Chen and D.A. Graetz. 1971. Importance of denitrification and nitrate reduction in sediments to the nitrogen budgets of lakes. *Nature* **233**: 66-67.
- Kelly, J. R. and B. L. Nowicki. 1992. Sediment denitrification in Boston Harbor. Massachusetts Water Resource Authority. Environmental Quality Department Technical Report 92-2. 31 p.
- . 1993. Direct measurements of denitrification in Boston Harbor and Massachusetts Bay sediments. Massachusetts Water Resource Authority. Environmental Quality Department Technical Report 93-3. 21 p.
- Kemp, W. M., P.Sampou, J.Caffrey, M.Mayer, K. Henriksen, and W.R. Boynton. 1990. Ammonium recycling versus denitrification in Chesapeake Bay sediments. *Limnology & Oceanography* **35**: 1545-1563.
- Koch, M. S., E. Maltby, G.A. Oliver and S.A. Bakker. 1992. Factors controlling denitrification rates of tidal mudflats and fringing salt marshes in south-west England. *Estuarine, Coastal and Shelf Science* **34**: 471-485.
- Koerselman, W., H. De Caluwe and W.M. Kiese kamp. 1989. Denitrification and dinitrogen fixation in two quaking fens in the Vechtplassen area, the Netherlands. *Biogeochemistry* **8**: 153-165.
- Koike, I. and A. Hattori. 1979. Estimates of denitrification in sediments of the Bering Sea shelf. *Deep-Sea Research* **26**: 409-415.
- Lamontagne, M. G., V. Astorga, E. Giblin and I. Valiela. 2002. Denitrification and the stoichiometry of nutrient regeneration in Waquoit Bay, Massachusetts. *Estuaries* **25**: 272-281.
- Lamontagne, M. G. and I. Valiela. 1995. Denitrification measured by a direct N sub(2) flux method in sediments of Waquoit Bay, MA. *Biogeochemistry* **31**: 63-83.
- Law, C. S., A.P. Rees and N.P. Owens. 1991. Temporal variability of denitrification in estuarine sediments. *Estuarine, Coastal and Shelf Science* **33**: 37-56.
- Likens, G. E. and O. L. Loucks. 1978. Analysis of five North American lake ecosystems. 3. Sources, loading and fate of nitrogen and phosphorus. *Int. Ver. Theor. Angew. Limnol. Verh.* **20**: 568-573.

- Lindau, C. W. and R. D. Delaune. 1991. Dinitrogen and nitrous oxide emission and entrapment in *Spartina alterniflora* saltmarsh soils following addition of N-15 labelled ammonium and nitrate. *Estuarine, Coastal and Shelf Science* **32**: 161-172.
- Lohse, L., H.T. Kloosterhuis, W. Van Raaphorst and W. Helder. 1996. Denitrification rates as measured by the isotope pairing method and by the acetylene inhibition technique in continental shelf sediments of the North Sea. *Marine Ecology Progress Series* **132**: 169-179.
- Lowrance, R. 1992. Nitrogen outputs from a field-size agricultural watershed. *Journal of Environmental Quality* **21**: 602-607.
- Lowrance, R., G. Vellidis and R.K. Hubbard. 1995. Denitrification in a restored riparian forest wetland. *Journal of Environmental Quality* **24**: 808-815.
- Madsen, P. P. 1979. Seasonal variation of denitrification rate in sediment determined by use of ¹⁵N. *Water Research* **13**: 461-465.
- Mengis, M., R. Gachter, B. Wehrli and S. Bernasconi. 1997. Nitrogen elimination in two deep eutrophic lakes. *Limnology & Oceanography* **42**: 1530-1543.
- Merrill, A. G. and D. R. Zak. 1992. Factors controlling denitrification rates in upland and swamp forests. *Can. J. For. Res.* **22**: 1597-1604.
- Messer, J. and P. L. Brezonik. 1983. Comparison of denitrification rate estimation techniques in a large, shallow lake. *Water Research* **17**: 631-640.
- Michotey, V. and P. Bonin. 1997. Evidence for anaerobic bacterial processes in the water column: Denitrification and dissimilatory nitrate ammonification in the northwestern Mediterranean Sea. *Marine Ecology Progress Series* **160**: 47-56.
- Miyajima, T., M. Suzumura, Y. Umezawa and I. Koike. 2001. Nitrogen transformation in carbonate sediments of a coral-reef lagoon and associated seagrass beds. *Marine Ecology Progress Series* **217**: 273-286.
- Morell, J. M. and J. E. Corredor. 1993. Sediment nitrogen trapping in a mangrove lagoon. *Estuarine, Coastal and Shelf Science* **37**: 203-212.
- Nedwell, D. B., T.H. Blackburn and W.J. Wiebe. 1994. Dynamic nature of the turnover of organic carbon, nitrogen and sulphur in the sediments of a Jamaican mangrove forest. *Marine Ecology Progress Series* **110**: 223-231.

- Nielsen, K., L.P. Nielsen and P. Rasmussen. 1995. Estuarine nitrogen retention independently estimated by the denitrification rate and mass balance methods: a study of Norsminde Fjord, Denmark. *Marine Ecology Progress Series* **119**: 275-283.
- Nielsen, K., N. Risgaard-Petersen, B. Soemod, S. Rysgaard and T. Bergoe. 2001. Nitrogen and phosphorus retention estimated independently by flux measurements and dynamic modelling in the estuary, Randers Fjord, Denmark. *Marine Ecology Progress Series* **219**: 25-40.
- Nielsen, L. P., P.B. Christensen, N.P. Revsbech and J. Sorensen. 1990. Denitrification and photosynthesis in stream sediment studied with microsensor and whole-core techniques. *Limnology & Oceanography* **35**: 1135-1144.
- Nielsen, L. P. 1992. Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiology Ecology* **86**: 357-362.
- Nielsen, L. P. and R. Glud. 1996. Denitrification in a coastal sediment measured in situ by the nitrogen isotope pairing technique applied to a benthic flux chamber. *Marine Ecology Progress Series* **137**: 181-186.
- Nishio, T., I. Koike and A. Hatorri. 1982. Denitrification, nitrate reduction and oxygen consumption in coastal and estuarine sediments. *Applied and Environmental Microbiology* **43**: 648-653.
- . 1983. Estimates of denitrification and nitrification in coastal and estuarine sediments. *Applied and Environmental Microbiology* **45**: 445-450.
- Nowicki, B. L. 1994. The effect of temperature, oxygen, salinity, and nutrient enrichment on estuarine denitrification rates measured with a modified nitrogen gas flux technique. *Estuarine, Coastal and Shelf Science* **38**: 137-156.
- Nowicki, B. L., J.R. Kelly, E. Requentina and D. Van Keuren. 1997. Nitrogen losses through sediment denitrification in Boston Harbor and Massachusetts Bay. *Estuaries* **20**: 626-639.
- Nowicki, B. L., E. Requentana, D. Van Keuren and J. Portnoy. 1999. The role of sediment denitrification in reducing groundwater-derived nitrate inputs to Nauset Marsh Estuary, Cape Cod, Massachusetts. *Estuaries* **22**: 245-259.
- Ogilvie, B., D.B. Nedwell, R.M. Harrison, A. Robinson, and A. Sage. 1997. High nitrate, muddy estuaries as nitrogen sinks: the nitrogen budget of the River Colne estuary (United Kingdom). *Marine Ecology Progress Series* **150**: 217-228.

- Oremland, R. S., C. Umberger, C.W. Culbertson and R.L. Smith. 1984. Denitrification in San Francisco Bay intertidal sediments. *Applied and Environmental Microbiology* **47**: 1106-1112.
- Oren, A. and T. H. Blackburn. 1979. Estimation of sediment denitrification rates at in situ nitrate concentrations. *Applied and Environmental Microbiology* **37**: 174-176.
- Ottosen, L. D., N. Risgaard-Petersen, L.P. Nielsen and T. Dalsgaard. 2001. Denitrification in exposed intertidal mud-flats, measured with a new ¹⁵N-ammonium spray technique. *Marine Ecology Progress Series* **209**: 35-42.
- Patrick, W. H. and K. R. Reddy. 1976. Denitrification reactions in flooded soils and water bottoms: dependence on oxygen supply and ammonium diffusion. *Journal of Environmental Quality* **4**: 469-471.
- Pelegri, S. P., L.P. Nielsen and T.H. Blackburn. 1994. Denitrification in estuarine sediment stimulated by the irrigation activity of the amphipod *Corophium volutator*. *Marine Ecology Progress Series* **105**: 285-290.
- Reddy, K. R., W.H. Patrick Jr. And C.W. Lindau. 1989. Nitrification-denitrification at the plant root-sediment interface in wetlands. *Limnology & Oceanography* **34**: 1004-1013.
- Richards, F. A. and W. W. Broenkow. 1971. Chemical changes, including nitrate reduction, in Darwin Bay, Galapagos archipelago over a 2-month period, 1969. *Limnology & Oceanography* **16**: 758-765.
- Risgaard-Petersen, N., S. Rysgaard, L.P. Nielsen and N.P. Revsbech. 1994. Diurnal variation of denitrification and nitrification in sediments colonized by benthic microphytes. *Limnology & Oceanography* **39**: 573-579.
- Risgaard-Petersen, N., T. Dalsgaard, S. Rysgaard, P.B. Christensen, J. Borum, J; K. Mcglathery and L.P. Nielsen. 1998. Nitrogen balance of a temperate eelgrass *Zostera marina* bed. *Marine Ecology Progress Series* **174**: 281-191.
- Risgaard-Petersen, N. and K. Jensen. 1997. Nitrification and denitrification in the rhizosphere of the aquatic macrophyte *Lobelia dortmanna* L. *Limnology & Oceanography* **42**: 529-537.
- Risgaard-Petersen, N. and L. D. Ottosen. 2000. Nitrogen cycling in two temperate *Zostera marina* beds: seasonal variation. *Marine Ecology Progress Series* **198**: 93-107.
- Rivera-Monroy, V. and R. R. Twilley. 1996. The relative role of denitrification and immobilization in the fate of inorganic nitrogen in mangrove sediments (Terminos Lagoon, Mexico). *Limnology & Oceanography* **41**: 284-296.

- Rivera-Monroy, V. H., R.R. Twilley, R.G. Boustany, J.W. Day, F. Vera-Herrera and M. Del Carmen Ramirez. 1995. Direct denitrification in mangrove sediments in Terminos Lagoon, Mexico. *Marine Ecology Progress Series* **126**: 97-109.
- Robinson, J. B., H.R. Whiteley, W. Stammers, N. Kaushik and P. Sain. 1979. The fate of nitrate in small streams and its management implications, p. 247-259. *In Proc. 10th Annu. Agric. Waste Manage. Conf. Cornell University.*
- Ronner, U. and F. Sorensson. 1985. Denitrification rates in the low-oxygen waters of the stratified Baltic proper. *Applied and Environmental Microbiology* **50**: 801-806.
- Rudd, J. W., C.A. Kelly, V. St. Louis, R.H. Hesslein, A. Furutani and M.H. Holoka. 1986. Microbial consumption of nitric and sulfuric acids in acidified north temperate lakes. *Limnology & Oceanography* **31**: 1267-1280.
- Rudd, J. W., C.A. Kelly, D.W. Schindler, D.W. and M.A. Turner. 1990. A comparison of the acidification efficiencies of nitric and sulfuric acids by two whole-lake addition experiments. *Limnology & Oceanography* **35**: 663-679.
- Rysgaard, S., N. Risgaard-Petersen, N.P. Sloth, K. Jensen and L.P. Nielsen. 1994. Oxygen regulation of nitrification and denitrification in sediments. *Limnology & Oceanography* **39**: 1643-1652.
- Rysgaard, S., P.B. Christensen and L.P. Nielsen. 1995. Seasonal variation in nitrification and denitrification in estuarine sediment colonized by benthic microalgae and bioturbating infauna. *Marine Ecology Progress Series* **126**: 111-121.
- Rysgaard, S., B. Thamdrup, N. Risgaard-Petersen, H. Fossing, P. Berg, P.B. Christensen, and T. Dalsgaard. 1998. Seasonal carbon and nutrient mineralization in a high-Arctic coastal marine sediment, Young Sound, Northeast Greenland. *Marine Ecology Progress Series* **175**: 261-276.
- Rysgaard, S., P. Thatstum, T. Dalsgaard, P.B. Christensen and N.P. Sloth. 1999. Effects of salinity on NH₄ adsorption capacity, nitrification and denitrification in Danish estuarine sediments. *Estuaries* **22**: 21-30.
- Seitzinger, S. P., S.W. Nixon and M. Pilson. 1984. Denitrification and nitrous oxide production in a coastal marine ecosystem. *Limnology & Oceanography* **29**: 73-83.
- Seitzinger, S. P. 1987. Nitrogen biogeochemistry in an unpolluted estuary: the importance of benthic denitrification. *Marine ecology progress series* **41**: 177-186.
- . 1988. Denitrification in freshwater and coastal marine ecosystems: ecological and geochemical significance. *Limnology & Oceanography* **33**: 702-724.

- Seitzinger, S. P., L.P. Nielsen, J. Caffrey and P.B. Christensen. 1993. Denitrification measurements in aquatic sediments: a comparison of three methods. *Biogeochemistry* **23**: 147-167.
- Seitzinger, S. P. 1994. Linkages between organic matter mineralization and denitrification in eight riparian wetlands. *Biogeochemistry* **25**: 19-39.
- Seitzinger, S. P. and A. E. Giblin. 1996. Estimating denitrification in North Atlantic continental shelf sediments. *Biogeochemistry* **35**: 235–260.
- Seitzinger, S. P. and S. W. Nixon. 1985. Eutrophication and the rate of denitrification and N₂O production in coastal marine sediments. *Limnology & Oceanography* **30**: 1332-1339.
- Serruya, C. 1975. Nitrogen and phosphorus balances and load-biomass relationship in Lake Kinneret (Israel). *Int. Ver. Theor. Angew. Limnol. Verh.* **19**: 1357-1369.
- Shieh, W. Y. and C. M. Liu. 1996. Denitrification by a novel halophilic fermentative bacterium. *Canadian Journal of Microbiology* **42**: 507-514.
- Slater, J. M. and D. G. Capone. 1987. Denitrification in aquifer soils and nearshore marine sediments influenced by groundwater nitrate. *Applied and Environmental Microbiology* **53**: 1292-1297.
- Sloth, N. P., B. Riemann, L.P. Nielsen, Lp and T.H. Blackburn. 1996. Resilience of pelagic and benthic microbial communities to sediment resuspension in a coastal ecosystem, Knebel Vig, Denmark. *Estuarine, Coastal and Shelf Science* **42**: 405-415.
- Smith, C. J., R.D. Delaune and W.H. Patrick, Jr. 1985. Fate of riverine nitrate entering an estuary: I. Denitrification and nitrogen burial. *Estuaries* **8**: 15-21.
- Sorensen, J. 1978. Denitrification rates in a marine sediment as measured by the acetylene inhibition technique. *Applied and Environmental Microbiology* **36**: 139-143.
- Sorensen, J., B.B. Jorgensen and N.P. Revsbech. 1979. A comparison of oxygen, nitrate and sulfate respiration in coastal marine sediments. *Marine Ecology* **5**: 105-115.
- Sorensen, J., K. Jorgensen, S. Colley, D. Hydes, J. Thomson and T. Wilson. 1987. Depth localization of denitrification in a deep-sea sediment from the Madeira Abyssal Plain. *Limnology & Oceanography* **32**: 758-762.
- Stockenberg, A. and R. W. Johnstone. 1997. Benthic denitrification in the Gulf of Bothnia. *Estuarine, Coastal and Shelf Science* **45**: 835-843.

- Sundbäck, K., A. Miles and E. Göransson. 2000. Nitrogen fluxes, denitrification and the role of microphytobenthos in microtidal shallow-water sediments: an annual study. *Marine Ecology Progress Series* **2000**: 59-76.
- Svensson, J. M., G. Carrer and M. Bocci. 2000. Nitrogen cycling in sediments of the Lagoon of Venice, Italy. *Marine Ecology Progress Series* **199**: 1-11.
- Thompson, S. P., H.W. Paerl and M.C. Go. 1995. Seasonal patterns of nitrification and denitrification in a natural and a restored salt marsh. *Estuaries* **18**: 399-408.
- Trimmer, M., D.B. Nedwell, D.B. Sivyer and S.J. Malcolm. 1998. Nitrogen fluxes through the lower estuary of the River Great Ouse, England: the role of the bottom sediments. *Marine Ecology Progress Series* **163**: 109-124.
- Trimmer, M., R.J. Gowen, B.M. Stewart and D.B. Nedwell. 1999. The spring bloom and its impact on benthic mineralisation rates in western Irish Sea sediments. *Marine Ecology Progress Series* **185**: 37-46.
- Trimmer, M., D.B. Nedwell, D.B. Sivyer and S.J. Malcolm. 2000a. Seasonal benthic organic matter mineralisation measured by oxygen uptake and denitrification along a transect of the inner and outer River Thames estuary, UK. *Marine Ecology Progress Series* **197**: 103-119.
- . 2000b. Seasonal organic mineralisation and denitrification in intertidal sediments and their relationship to the abundance of *Enteromorpha* sp. and *Ulva* sp. *Marine Ecology Progress Series* **203**: 67-80.
- Tuominen, L., A. Heinaenen, J. Kuparinen and L.P. Nielsen. 1998. Spatial and temporal variability of denitrification in the sediments of the northern Baltic Proper. *Marine Ecology Progress Series* **172**: 13-24.
- Tuominen, L., K. Maekelae, K. Lehtonen, H. Haahti, S. Hietanen and J. Kuparinen. 1999. Nutrient fluxes, porewater profiles and denitrification in sediment influenced by algal sedimentation and bioturbation by *Monoporeia affinis*. *Estuarine, Coastal and Shelf Science* **49**: 83-97.
- Twilley, R. and W. M. Kemp. 1987. Estimates of sediment denitrification and its influence on the fate of nitrogen in Chesapeake Bay. US Environmental Protection Agency, Chesapeake Bay Program. 47 p.
- Urban, N. R., S.J. Eisenreich and S.E. Bayley. 1988. The relative importance of denitrification and nitrate assimilation in midcontinental bogs. *Limnology & Oceanography* **33**: 1611-1617.

- Usui, T., I. Koike and N. Ogura. 2001. N₂O Production, Nitrification and Denitrification in an Estuarine Sediment. *Estuarine, Coastal and Shelf Science* **52**: 769-781.
- Valiela, I., M.L. Cole, J. McClelland, J. Hauxwell and J. Cebrian. 2000. Role of salt marshes as part of coastal landscapes, p. 23-38. *In* M. P. W. a. P. A. Kreeger [ed.], *Concepts and Controversies in Tidal Marsh Ecology*. Kluwer Academic Publishers.
- Vanderborght, J. and G. Billen. 1975. Vertical distribution of nitrate concentration in interstitial water of marine sediments with nitrification and denitrification. *Limnology & Oceanography* **22**: 794-803.
- Wang, F., S.K. Juniper, S.P. Pelegri and S.A. Macko. 2003. Denitrification in sediments of the Laurentian Trough, St. Lawrence Estuary, Quebec, Canada. *Estuarine, Coastal and Shelf Science* **57**: 515-522.
- Welsh, D., M. Bartoli, D. Nizzoli, G. Castaldelli, S.A. Riou and P. Viaroli. 2000. Denitrification, nitrogen fixation, community primary productivity and inorganic-N and oxygen fluxes in an intertidal *Zostera noltii* meadow. *Marine Ecology Progress Series* **208**: 65-77.
- Westermann, P. and B. K. Ahring. 1987. Dynamics of methane production, sulfate reduction and denitrification in a permanently waterlogged alder swamp. *Applied and Environmental Microbiology* **53**: 2554-2559.
- White, D. S. 1994. ¹⁵N₂ studies of nitrogen retention in the vegetated sediments of a New England salt marsh, Thesis. Boston University, Boston Massachusetts, USA.
- White, D. S. and B. L. Howes. 1994. Long-term super(¹⁵N)-nitrogen retention in the vegetated sediments of a New England salt marsh. *Limnology & Oceanography* **39**: 1878-1892.
- Xue, Y., D. A. Kovacic, M. B. David, L. E. Gentry, R. L. Mulvaney, and C. W. Lindau. 1999. In situ measurements of denitrification in constructed wetlands. *Journal of Environmental Quality* **28**: 263-269.
- Yoon, W. B. and R. Benner. 1992. Denitrification and oxygen consumption in sediments of two south Texas estuaries. *Marine Ecology Progress Series* **90**: 157-167.
- Zak, D. R. and D. F. Grigal. 1991. Nitrogen mineralization, nitrification and denitrification in upland and wetland ecosystems. *Oecologia* **88**: 189-196.
- Zimmerman, A. and R. Benner. 1994. Denitrification, nutrient regeneration and carbon mineralization in sediments of Galveston Bay, Texas, USA. *Marine Ecology Progress Series* **114**: 275-288.